



## **Bridging the gap between Transcriptome and Metabolome by using Genome-scale Metabolic Models**

**Zelezniak, Aleksej; Patil, Kiran Raosaheb**

*Publication date:*  
2009

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Zelezniak, A., & Patil, K. R. (2009). *Bridging the gap between Transcriptome and Metabolome by using Genome-scale Metabolic Models*. Poster session presented at 1st Systems Biology as a driver for Industrial Biotechnology Workshop, Istanbul, Turkey. <http://www.sysbio.se/SYSINBIO/index.html>

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



## Motivation


The direct *in vivo* investigation of metabolism is complicated by the fact that it is not yet possible to quantitatively measure all metabolite levels within the cell. Thus, there is a need for development of methods that enable to predict whether a particular metabolite is being accumulated or depleted following a genetic/environmental perturbation. Towards this goal, a computational method is presented for integrating transcriptome data and genome-scale metabolic models for predicting accumulation or depletion of intracellular metabolites.

## Methods

### Reaction Directionality Analysis (RDA)


#### Subject to

Fixed fermentation physiological parameters, network stoichiometry



Linear programming (LP) problem is solved for maximizing and minimizing each flux  $f_i$ , thus resulting in a system flexibility under constraint of fixed substrates uptake and biomass production. In other words, the method aims at checking each reaction's behavior – whether it can go either in both directions but still maintaining main flux towards cellular objective production, or only in one direction.

### Metabolite Scoring System



$$S = \frac{\sum_{i=1}^j RP_i}{j} + \frac{\sum_{i=1}^k RC_i}{k}$$

$$RP_i = FP_i \times PP_i$$

$$RC_i = FC_i \times PC_i$$

$$FP_i = \begin{cases} 1 & \text{if } \log(\text{fold change}) \text{ of gene controlling } RP_i > 0 \\ -1 & \text{if } \log(\text{fold change}) \text{ of gene controlling } RP_i < 0 \end{cases}$$

$$PP_i = \begin{cases} 1 & \text{if } p\text{-value of gene controlling } RP_i \leq 0.05 \\ 0.5 & \text{if } p\text{-value of gene controlling } RP_i > 0.05 \end{cases}$$


$$FC_i = \begin{cases} -1 & \text{if } \log(\text{fold change}) \text{ of gene controlling } RC_i > 0 \\ 1 & \text{if } \log(\text{fold change}) \text{ of gene controlling } RC_i < 0 \end{cases}$$

$$PC_i = \begin{cases} 1 & \text{if } p\text{-value of gene controlling } RC_i \leq 0.05 \\ 0.5 & \text{if } p\text{-value of gene controlling } RC_i > 0.05 \end{cases}$$

## Conclusions

The proposed method is able to qualitatively predict intracellular metabolite pool change between two different physiological conditions. Accumulation/depletion of metabolites were successfully predicted in 73% of the cases. Methodology can be used, for example, as a guiding tool for making targeted metabolome analysis and biomarker development. An interesting application will be in human diseases research, where transcriptomic phenotype is more readily measurable than metabolome.

## Results



- Out of 1433 metabolic reactions, 507 are predicted by RDA to have *forward* direction, *40 backward*, 230 (out of 405) *reversible* and 656 *infeasible*.
- Changes in 220 metabolite pools are predicted, out of them - 112 are being accumulated, 107 depleted and 49 are at homeostasis state
- Experimental measurements of 11 metabolites levels [1] were used as a proof of concept, the direction of change of 8 metabolites were correctly predicted.

#### References:

- Kresnowati MT, van Winden WA, Almering MJ, ten Pierick A, Ras C, et al. (2006) When transcriptome meets metabolome: fast cellular responses of yeast to sudden relief of glucose limitation. *Mol Syst Biol* 2: 49.